



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 3/37	A1	(11) International Publication Number: WO 99/27053 (43) International Publication Date: 3 June 1999 (03.06.99)
(21) International Application Number: PCT/US98/24699 (22) International Filing Date: 20 November 1998 (20.11.98) (30) Priority Data: 60/066,346 21 November 1997 (21.11.97) US 60/087,640 2 June 1998 (02.06.98) US (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KASTURI, Chandrika [US/US]; 10044 Cliffwood Court, Cincinnati, OH 45241 (US). SCHAFER, Michael, Gayle [US/US]; 6032 South Kramer Drive, Alexandria, KY 41001 (US). SIVIK, Mark, Robert [US/US]; Apartment F, 40 Clubhouse Lane, Fairfield, OH 45014 (US). KLUESENER, Bernard, William [US/US]; 208 Circle Drive, Harrison, OH 45030 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).	(81) Designated States: CN, CZ, CZ (Utility model), JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: FOAM STABLE LIQUID DISHWASHING COMPOSITIONS (57) Abstract The present invention relates to zwitterionic polymer suds stabilizers suitable for use in liquid detergent compositions for hand dishwashing. A 10 % aqueous solution of this composition has a pH from 4 to 12.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

FOAM STABLE LIQUID DISHWASHING COMPOSITIONS

FIELD OF THE INVENTION

The present invention relates to liquid detergent compositions useful for hand dishwashing, said compositions comprising a zwitterionic polymer suds stabilizer which provides increased suds volume and suds duration. The present invention also relates to methods for providing increased suds volume and suds duration during cleaning of dishware and other hand washing operations.

BACKGROUND OF THE INVENTION

Liquid detergent compositions which are suitable for hand dishwashing must satisfy several criteria in order to be effective. These compositions must be effective in cutting grease and greasy food material and once removed, must keep the greasy material from re-depositing on the dishware.

The presence of suds in a hand dishwashing operation has long been used as a signal that the detergent continues to be effective. However, depending upon the circumstances, the presence of suds or the lack thereof, has no bearing upon the efficacy of liquid detergents. Therefore, the consumer has come to rely upon a somewhat erroneous signal, the lack or absence of soap suds, to indicate the need for additional detergent. In many instances the consumer is adding an additional amount of detergent far in excess of the amount necessary to thoroughly clean the dishes. This wasteful use of detergent is especially true in hand dishwashing since the soiled cooking articles are usually cleaned in a "washing difficulty" queue, for example, glasses and cups, which usually do not contact greasy food, are washed first, followed by plates and flatware, and finally pots and pans which contain the most residual food material and are usually, therefore, the "greasiest".

The lack of suds in the dishwater when pots and pans are usually cleaned, together with the visual inspection of the amount of residual food material on the cookware surface, typically compels the consumer to add additional detergent when a sufficient amount still remains in solution to effectively remove the soil and grease from the dishware or cookware surface. However, effective grease cutting materials do not necessarily produce a substantial amount of corresponding suds.

Accordingly, there remains a need in the art for liquid dishwashing detergents useful for hand washing dishware which have an enduring suds level while maintaining effective grease cutting properties. The need exists for a composition which can maintain a high level of suds as long as the dishwashing composition is effective.

Indeed, there is a long felt need to provide a hand dishwashing composition which can be use efficiently by the consumer such that the consumer uses only the necessary amount of detergent to fully accomplish the cleaning task.

SUMMARY OF THE INVENTION

The present invention meets the aforementioned needs in that it has been surprisingly discovered that certain zwitterionic polymers serve as suds extenders and suds volume enhancers in liquid detergent compositions which are effective in cutting grease.

The zwitterionic polymeric suds stabilizers of the present invention comprise monomeric units which have at least one moiety capable of sustaining a negative charge at a pH of from about 4 to about 12 and at least one moiety capable of sustaining a positive charge within the same pH range.

A first aspect of the present invention relates to detergent compositions suitable for use in hand dishwashing, said compositions comprising:

- a) an effective amount of a zwitterionic polymeric suds stabilizer;
- b) an effective amount of a deterative surfactant; and
- c) the balance carriers and other adjunct ingredients;

provided the pH of a 10% aqueous solution of said composition is from about 4 to about 12.

The present invention further relates to zwitterionic polymeric materials which are suitable for use in detergents wherein the formulator desires to extend the amount and duration of suds.

The present invention also relates to methods for providing increased suds and increased duration of suds while hand washing dishware comprising the step of dissolving a composition according to the present invention in water to form a hand dish washing solution and then washing dishware by hand in said solution. These and other objects, features and advantages will become apparent to those of ordinary skill in the art from a reading of the following detailed description and the appended claims.

All percentages, ratios and proportions herein are by weight, unless otherwise specified. All temperatures are in degrees Celsius (°C) unless otherwise specified. All documents cited are in relevant part, incorporated herein by reference.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to liquid detergent compositions suitable for hand washing, especially for hand washing dishware and cookware, said compositions comprising a zwitterionic polymer which serves to provide enhanced suds volume and suds duration. The zwitterionic polymers of the present invention may be homopolymers

or copolymers, each of which may be suitably crosslinked. The zwitterionic polymers are comprised of moieties which when placed in an aqueous solution having a pH of from 4 to about 12, said moieties are capable of sustaining a positive or negative charge.


The detergent compositions of the present invention have a pH of from about 4 to about 12 when measured as a 10% aqueous solution. The polymeric suds enhancers of the present invention are zwitterionic polymers. For the purposes of the present invention the term "zwitterionic polymer" is defined as "a polymeric material comprised of one or more monomers wherein each monomer has one or more moieties capable of sustaining a positive or negative charge at a pH of from about 4 to about 12 such that the number of positively charged moieties is equal to the number of negatively charged moieties at the isoelectric point of said polymer."

The compositions according to the present invention also comprises an effective amount of one or more deterative surfactants described herein below as well as carriers and other adjunct ingredients. The polymers and copolymers of the present invention act together with said surfactants and other adjunct ingredients, especially certain diamines, to provide for efficient grease cutting and anti-redeposition of grease.

Zwitterionic Polymers

The polymeric suds stabilizers of the present invention are homopolymers or copolymers wherein the monomers which comprise said homopolymers or copolymers contain a moiety capable of being protonated at a pH of from about 4 to about 12, or a moiety capable of being de-protonated at a pH of from about 4 to about 12, of a mixture of both types of moieties.

A preferred class of zwitterionic polymer suitable for use as a suds volume and suds duration enhancer has the formula:




wherein R is C₁-C₁₂ linear alkylene, C₁-C₁₂ branched alkylene, and mixtures thereof; preferably C₁-C₄ linear alkylene, C₃-C₄ branched alkylene; more preferably methylene and 1,2-propylene. R¹ and R² are defined herein after. The index x is from 0 to 6; y is 0 or 1; z is 0 or 1.

The index n has the value such that the zwitterionic polymers of the present invention have an average molecular weight of from about 1,000 to about 2,000,000 preferably from about 5,000 to about 1,000,000, more preferably from about 10,000 to about 750,000, more preferably from about 20,000 to about 500,000, even


more preferably from about 35,000 to about 300,000 daltons. The molecular weight of the polymeric suds boosters, can be determined via conventional gel permeation chromatography.

Anionic Units


R^1 is a unit capable of having a negative charge at a pH of from about 4 to about 12. Preferred R^1 has the formula:




wherein L is a linking unit independently selected from the following:



mixtures thereof, wherein R' is independently hydrogen, C_1 - C_4 alkyl, and mixtures thereof; preferably hydrogen or alternatively R' and S can form a heterocycle of 4 to 7 carbon atoms, optionally containing other hetero atoms and optionally substituted. Preferably the linking group L can be introduced into the molecule as part of the original monomer backbone, for example, a polymer having L units of the formula:



can suitably have this moiety introduced into the polymer via a carboxylate containing monomer, for example, a monomer having the general formula:



When the index i is 0, L is absent.

For anionic units S is a "spacing unit" wherein each S unit is independently selected from C_1 - C_{12} linear alkylene, C_1 - C_{12} branched alkylene, C_3 - C_{12} linear alkenylene, C_3 - C_{12} branched alkenylene, C_3 - C_{12} hydroxyalkylene, C_4 - C_{12} dihydroxyalkylene, C_6 - C_{10} arylene, C_8 - C_{12} dialkylarylene, $-(R^5O)_kR^5-$, $-(R^5O)_kR^6(OR^5)_k-$, $-CH_2CH(OR^7)CH_2-$, and mixtures thereof; wherein R^5 is C_2 - C_4 linear alkylene, C_3 - C_4 branched alkylene, and mixtures thereof, preferably ethylene, 1,2-propylene, and mixtures thereof, more preferably ethylene; R^6 is C_2 - C_{12} linear alkylene,

and mixtures thereof, preferably ethylene; R^7 is hydrogen, C_1 - C_4 alkyl, and mixtures thereof, preferably hydrogen. The index k is from 1 to about 20.

Preferably S is C_1 - C_{12} linear alkylene, $-(R^5O)_kR^5$ -, and mixtures thereof. When S is a $-(R^5O)_kR^5$ - unit, said units may be suitably formed by the addition of an alkyleneoxy producing reactant (e.g. ethylene oxide, epichlorohydrin) or by addition of a suitable polyethyleneglycol. More preferably S is C_2 - C_4 linear alkylene. When the index j is 0 the S unit is absent.


R^3 is independently selected from hydrogen, $-CO_2M$, $-SO_3M$, $-OSO_3M$, $-CH_2P(O)(OM)_2$, $-OP(O)(OM)_2$, units having the formula:




wherein each R^8 , R^9 , and R^{10} is independently selected from the group consisting of hydrogen, $-(CH_2)_mR^{11}$, and mixtures thereof, wherein R^{11} is $-CO_2H$, $-SO_3M$, $-OSO_3M$, $-CH(CO_2H)CH_2CO_2H$, $-CH_2P(O)(OH)_2$, $-OP(O)(OH)_2$, and mixtures thereof, preferably $-CO_2H$, $-CH(CO_2H)CH_2CO_2H$, and mixtures thereof, more preferably $-CO_2H$; provided that one R^8 , R^9 , or R^{10} is not a hydrogen atom, preferably two R^8 , R^9 , or R^{10} units are hydrogen. M is hydrogen or a salt forming cation, preferably hydrogen. The index m has the value from 0 to 10.

Cationic Units


R^2 is a unit capable of having a positive charge at a pH of from about 4 to about 12. Preferred R^2 has the formula:



wherein L^1 is a linking unit independently selected from the following:



and mixtures thereof; wherein R' is independently hydrogen, C₁-C₄ alkyl, and mixtures thereof; preferably hydrogen or alternatively R' and S can form a heterocycle of 4 to 7 carbon atoms, optionally containing other hetero atoms and optionally substituted. Preferably L¹ has the formula:




When the index i' is equal to 0, L¹ is absent.

For cationic units S is a "spacing unit" wherein each S unit is independently selected from C₁-C₁₂ linear alkylene, C₁-C₁₂ branched alkylene, C₃-C₁₂ linear alkenylene, C₃-C₁₂ branched alkenylene, C₃-C₁₂ hydroxyalkylene, C₄-C₁₂ dihydroxyalkylene, C₆-C₁₀ arylene, C₈-C₁₂ dialkylarylene, -(R⁵O)_kR⁵-, -(R⁵O)_kR⁶(OR⁵)_k-, -CH₂CH(OR⁷)CH₂-, and mixtures thereof; wherein R⁵ is C₂-C₄ linear alkylene, C₃-C₄ branched alkylene, and mixtures thereof, preferably ethylene, 1,2-propylene, and mixtures thereof, more preferably ethylene; R⁶ is C₂-C₁₂ linear alkylene, and mixtures thereof, preferably ethylene; R⁷ is hydrogen, C₁-C₄ alkyl, and mixtures thereof, preferably hydrogen. The index k is from 1 to about 20.

Preferably S is C₁-C₁₂ linear alkylene, and mixtures thereof. Preferably S is C₂-C₄ linear alkylene. When the index j' is 0 the S unit is absent.


R⁴ is independently selected from amino, alkylamino carboxamide, 3-imidazolyl, 4-imidazolyl, 2-imidazolyl, 4-imidazolyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, piperazinyl, 2-

pyrrolidinyl, 3-pyrrolidinyl, guanidino, amidino, and mixtures thereof, preferably dialkylamino having the formula:




wherein each R^{11} is independently hydrogen, C_1 - C_4 alkyl, and mixtures thereof, preferably hydrogen or methyl or alternatively the two R^{11} can form a heterocycle of 4 to 8 carbon atoms, optionally containing other hetero atoms and optionally substituted.

An example of a preferred zwitterionic polymer according to the present invention has the formula:




wherein X is C_6 , n has a value such that the average molecular weight is from about 5,000 to about 1,000,000 daltons.

Further preferred zwitterionic polymers according to the present invention are polymers comprising monomers wherein each monomer has only cationic units or anionic units, said polymers have the formula:




wherein R, R^1 , x, y, and z are the same as defined herein above; $n^1 + n^2 = n$ such that n has a value wherein the resulting zwitterionic polymer has a molecular weight of form about 5,000 to about 1,000,000 daltons.

An example of a polymer having monomers with only an anionic unit or a cationic unit has the formula:




wherein the sum of n^1 and n^2 provide a polymer with an average molecular weight of from about 5,000 to about 750,000 daltons.

Another preferred zwitterionic polymer according to the present invention are polymers which have limited crosslinking, said polymers having the formula:



wherein R, R¹, L¹, S, j', x, y, and z are the same as defined herein above; n' is equal to n", and the value n' + n" is less than or equal to 5% of the value of n¹ + n² = n; n provides a polymer with an average molecular weight of from about 1,000 to about 2,000,000 daltons. R¹² is nitrogen, C₁-C₁₂ linear alkylene amino alkylene having the formula:



L^1 , and mixtures thereof, wherein each R^{13} is independently L^1 or ethylene.

The zwitterionic polymers of the present invention may comprise any combination of monomer units, for example, several different monomers having various R¹ and R² groups can be combined to form a suitable suds stabilizer. Alternatively the same R¹ unit may be used with a selection of different R² units and vice versa.

The liquid detergent compositions according to the present invention comprise at least an effective amount of one or more zwitterionic polymeric suds stabilizers described herein, preferably from about 0.01% to about 10%, more preferably from about 0.05% to about 5%, most preferably from about 0.1% to about 2% by weight, of said composition. What is meant herein by "an effective amount of zwitterionic polymeric suds stabilizer" is that the suds produced by the presently described compositions are sustained for an increased amount of time relative to a composition which does not comprise a zwitterionic polymeric suds stabilizer described herein. Additionally, the polymeric suds stabilizer can be present as the free base or as a salt. Typical counter ions include, citrate, maleate, sulfate, chloride, etc.

Detersive Surfactants

Anionic Surfactants - The anionic surfactants useful in the present invention are preferably selected from the group consisting of, linear alkylbenzene sulfonate, alpha olefin sulfonate, paraffin sulfonates, alkyl ester sulfonates, alkyl sulfates, alkyl alkoxy sulfate, alkyl sulfonates, alkyl alkoxy carboxylate, alkyl alkoxyated sulfates, sarcosinates, taurinates, and mixtures thereof. An effective amount, typically from about 0.5% to about 90%, preferably about 5% to about 60%, more preferably from about 10 to about 30%, by weight of anionic detersive surfactant can be used in the present invention.

Alkyl sulfate surfactants are another type of anionic surfactant of importance for use herein. In addition to providing excellent overall cleaning ability when used in combination with polyhydroxy fatty acid amides (see below), including good grease/oil cleaning over a wide range of temperatures, wash concentrations, and wash times, dissolution of alkyl sulfates can be obtained, as well as improved formulability in liquid detergent formulations are water soluble salts or acids of the formula $ROSO_3M$ wherein R preferably is a C_{10} - C_{24} hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C_{10} - C_{20} alkyl component, more preferably a C_{12} - C_{18} alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali (Group IA) metal cation (e.g., sodium, potassium, lithium), substituted or unsubstituted ammonium cations such as methyl-, dimethyl-, and trimethyl ammonium and quaternary ammonium cations, e.g., tetramethyl-ammonium and dimethyl piperdinium, and cations derived from alkanolamines such as ethanolamine, diethanolamine, triethanolamine, and mixtures thereof, and the like. Typically, alkyl chains of C_{12} - C_{16} are preferred for lower wash temperatures (e.g., below about 50°C) and C_{16} - C_{18} alkyl chains are preferred for higher wash temperatures (e.g., above about 50°C).

Alkyl alkoxyated sulfate surfactants are another category of useful anionic surfactant. These surfactants are water soluble salts or acids typically of the formula $RO(A)_mSO_3M$ wherein R is an unsubstituted C_{10} - C_{24} alkyl or hydroxyalkyl group having a C_{10} - C_{24} alkyl component, preferably a C_{12} - C_{20} alkyl or hydroxyalkyl, more preferably C_{12} - C_{18} alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl-, trimethyl-ammonium and quaternary ammonium cations, such as tetramethyl-ammonium, dimethyl piperidinium and cations derived from alkanolamines, e.g.

monoethanolamine, diethanolamine, and triethanolamine, and mixtures thereof. Exemplary surfactants are C₁₂-C₁₈ alkyl polyethoxylate (1.0) sulfate, C₁₂-C₁₈ alkyl polyethoxylate (2.25) sulfate, C₁₂-C₁₈ alkyl polyethoxylate (3.0) sulfate, and C₁₂-C₁₈ alkyl polyethoxylate (4.0) sulfate wherein M is conveniently selected from sodium and potassium. Surfactants for use herein can be made from natural or synthetic alcohol feedstocks. Chain lengths represent average hydrocarbon distributions, including branching.


Examples of suitable anionic surfactants are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23.

Secondary Surfactants - Secondary deterative surfactant can be selected from the group consisting of nonionics, cationics, ampholytics, zwitterionics, and mixtures thereof. By selecting the type and amount of deterative surfactant, along with other adjunct ingredients disclosed herein, the present detergent compositions can be formulated to be used in the context of laundry cleaning or in other different cleaning applications, particularly including dishwashing. The particular surfactants used can therefore vary widely depending upon the particular end-use envisioned. Suitable secondary surfactants are described below. Examples of suitable nonionic, cationic amphoteric and zwitterionic surfactants are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

Nonionic Detergent Surfactants - Suitable nonionic detergent surfactants are generally disclosed in U.S. Patent 3,929,678, Laughlin et al., issued December 30, 1975, at column 13, line 14 through column 16, line 6, incorporated herein by reference. Exemplary, non-limiting classes of useful nonionic surfactants include: amine oxides, alkyl ethoxylate, alkanoyl glucose amide, alkyl betaines, sulfobetaine and mixtures thereof.

Amine oxides are semi-polar nonionic surfactants and include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; water-soluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula



wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R^5 is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.


These amine oxide surfactants in particular include C_{10} - C_{18} alkyl dimethyl amine oxides and C_8 - C_{12} alkoxy ethyl dihydroxy ethyl amine oxides. Preferably the amine oxide is present in the composition in an effective amount, more preferably from about 0.1% to about 20%, even more preferably about 0.1% to about 15%, even more preferably still from about 0.5% to about 10%, by weight.

The polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols. In general, the polyethylene oxide condensates are preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 12 carbon atoms in either a straight chain or branched chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 5 to about 25 moles of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include Igepal® CO-630, marketed by the GAF Corporation; and Triton® X-45, X-114, X-100, and X-102, all marketed by the Rohm & Haas Company. These compounds are commonly referred to as alkyl phenol alkoxylates, (e.g., alkyl phenol ethoxylates).

The condensation products of aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Particularly preferred are the condensation products of alcohols having an alkyl group containing from about 10 to about 20 carbon atoms with from about 2 to about 18 moles of ethylene oxide per mole of alcohol. Examples of commercially available nonionic surfactants of this type include Tergitol® 15-S-9 (the condensation product of C_{11} - C_{15} linear secondary alcohol with 9 moles ethylene oxide), Tergitol® 24-L-6 NMW (the condensation product of C_{12} - C_{14} primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide


Corporation; Neodol® 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), Neodol® 23-6.5 (the condensation product of C₁₂-C₁₃ linear alcohol with 6.5 moles of ethylene oxide), Neodol® 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), Neodol® 45-4 (the condensation product of C₁₄-C₁₅ linear alcohol with 4 moles of ethylene oxide), marketed by Shell Chemical Company, and Kyro® EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company. Other commercially available nonionic surfactants include Dobanol 91-8® marketed by Shell Chemical Co. and Genapol UD-080® marketed by Hoechst. This category of nonionic surfactant is referred to generally as "alkyl ethoxylates."

The preferred alkylpolyglycosides have the formula



wherein R² is selected from the group consisting of alkyl, alkyl-phenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4- and/or 6-position, preferably predominantly the 2-position.

Fatty acid amide surfactants having the formula:




wherein R⁶ is an alkyl group containing from about 7 to about 21 (preferably from about 9 to about 17) carbon atoms and each R⁷ is selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, and -(C₂H₄O)_xH where x varies from about 1 to about 3.

Preferred amides are C₈-C₂₀ ammonia amides, monoethanolamides, diethanolamides, and isopropanolamides.

Preferably the nonionic surfactant, when present in the composition, is present in an effective amount, more preferably from about 0.1% to about 20%, even more preferably about 0.1% to about 15%, even more preferably still from about 0.5% to about 10%, by weight.

Polyhydroxy Fatty Acid Amide Surfactant - The detergent compositions hereof may also contain an effective amount of polyhydroxy fatty acid amide surfactant. By "effective amount" is meant that the formulator of the composition can select an amount of polyhydroxy fatty acid amide to be incorporated into the compositions that will improve the cleaning performance of the detergent composition. In general, for conventional levels, the incorporation of about 1%, by weight, polyhydroxy fatty acid amide will enhance cleaning performance.

The detergent compositions herein will typically comprise about 1% weight basis, polyhydroxy fatty acid amide surfactant, preferably from about 3% to about 30%, of the polyhydroxy fatty acid amide. The polyhydroxy fatty acid amide surfactant component comprises compounds of the structural formula:



wherein: R^1 is H, C_1 - C_4 hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl, or a mixture thereof, preferably C_1 - C_4 alkyl, more preferably C_1 or C_2 alkyl, most preferably C_1 alkyl (i.e., methyl); and R^2 is a C_5 - C_{31} hydrocarbyl, preferably straight chain C_7 - C_{19} alkyl or alkenyl, more preferably straight chain C_9 - C_{17} alkyl or alkenyl, most preferably straight chain C_{11} - C_{15} alkyl or alkenyl, or mixtures thereof; and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably will be derived from a reducing sugar in a reductive amination reaction; more preferably Z will be a glycityl. Suitable reducing sugars include glucose, fructose, maltose, lactose, galactose, mannose, and xylose. As raw materials, high dextrose corn syrup, high fructose corn syrup, and high maltose corn syrup can be utilized as well as the individual sugars listed above. These corn syrups may yield a mix of sugar components for Z. It should be understood that it is by no means intended to exclude other suitable raw materials. Z preferably will be selected from the group consisting of $-\text{CH}_2-(\text{CHOH})_n-\text{CH}_2\text{OH}$, $-\text{CH}(\text{CH}_2\text{OH})-(\text{CHOH})_{n-1}-\text{CH}_2\text{OH}$, $-\text{CH}_2-(\text{CHOH})_2(\text{CHOR}')(\text{CHOH})-\text{CH}_2\text{OH}$, and alkoxyated derivatives thereof, where n is an integer from 3 to 5, inclusive, and R' is H or a cyclic or aliphatic monosaccharide. Most preferred are glycityls wherein n is 4, particularly $-\text{CH}_2-(\text{CHOH})_4-\text{CH}_2\text{OH}$.

R' can be, for example, N-methyl, N-ethyl, N-propyl, N-isopropyl, N-butyl, N-2-hydroxy ethyl, or N-2-hydroxy propyl.

$R^2\text{-CO-N<}$ can be, for example, cocamide, stearamide, oleamide, lauramide, myristamide, capricamide, palmitamide, tallowamide, etc.


Z can be 1-deoxyglucityl, 2-deoxyfructityl, 1-deoxymaltityl, 1-deoxylactityl, 1-deoxygalactityl, 1-deoxymannityl, 1-deoxymaltotriosityl, etc.

Methods for making polyhydroxy fatty acid amides are known in the art. In general, they can be made by reacting an alkyl amine with a reducing sugar in a reductive amination reaction to form a corresponding N-alkyl polyhydroxyamine, and then reacting the N-alkyl polyhydroxyamine with a fatty aliphatic ester or triglyceride in a condensation/amidation step to form the N-alkyl, N-polyhydroxy fatty acid amide product. Processes for making compositions containing polyhydroxy fatty acid amides are disclosed, for example, in G.B. Patent Specification 809,060, published February 18, 1959, by Thomas Hedley & Co., Ltd., U.S. Patent 2,965,576, issued December 20, 1960 to E. R. Wilson, and U.S. Patent 2,703,798, Anthony M. Schwartz, issued March 8, 1955, and U.S. Patent 1,985,424, issued December 25, 1934 to Piggott, each of which is incorporated herein by reference.


Diamines

The preferred liquid detergent compositions of the present invention further comprise one or more diamines, preferably an amount of diamine such that the ratio of anionic surfactant present to the diamine is from about 40 : 1 to about 2 : 1. Said diamines provide for increased removal of grease and greasy food material while maintaining suitable levels of suds.

The diamines suitable for use in the compositions of the present invention have the formula:




wherein each R^{20} is independently selected from the group consisting of hydrogen, C_1 - C_4 linear or branched alkyl, alkyleneoxy having the formula:



wherein R^{21} is C_2 - C_4 linear or branched alkylene, and mixtures thereof; R^{22} is hydrogen, C_1 - C_4 alkyl, and mixtures thereof; y is from 1 to about 10; X is a unit selected from:

- i) C_3 - C_{10} linear alkylene, C_3 - C_{10} branched alkylene, C_3 - C_{10} cyclic alkylene, C_3 - C_{10} branched cyclic alkylene, an alkyleneoxyalkylene having the formula:



wherein R^{21} and y are the same as defined herein above;

- ii) C_3 - C_{10} linear, C_3 - C_{10} branched linear, C_3 - C_{10} cyclic, C_3 - C_{10} branched cyclic alkylene, C_6 - C_{10} arylene, wherein said unit comprises one or more electron donating or electron withdrawing moieties which provide said diamine with a pK_a greater than about 8; and

- iii) mixtures of (i) and (ii)

provided said diamine has a pK_a of at least about 8.


The preferred diamines of the present invention have a pK_1 and pK_2 which are each in the range of from about 8 to about 11.5, preferably in the range of from about 8.4 to about 11, more preferably from about 8.6 to about 10.75. For the purposes of the present invention the term " pK_a " stands equally well for the terms " pK_1 " and " pK_2 " either separately or collectively. The term pK_a as used herein throughout the present specification in the same manner as used by those of ordinary skill in the art. pK_a values are readily obtained from standard literature sources, for example, "Critical Stability Constants: Volume 2, Amines" by Smith and Martel, Plenum Press, N.Y. and London, (1975).

As an applied definition herein, the pK_a values of the diamines are specified as being measured in an aqueous solution at 25° C having an ionic strength of from about 0.1 to about 0.5 M. As used herein, the pK_a is an equilibrium constant dependent upon temperature and ionic strength, therefore, value reported by literature references, not measured in the above described manner, may not be within full agreement with the values and ranges which comprise the present invention. To eliminate ambiguity, the relevant conditions and/or references used for pK_a 's of this invention are as defined herein or in "Critical Stability Constants: Volume 2, Amines". One typical method of measurement is the potentiometric titration of the acid with sodium hydroxide and determination of the pK_a by suitable methods as described and referenced in "The Chemist's Ready Reference Handbook" by Shugar and Dean, McGraw Hill, NY, 1990.


Preferred diamines for performance and supply considerations are 1,3-bis(methylamino)cyclohexane, 1,3-diaminopropane ($pK_1=10.5$; $pK_2=8.8$), 1,6-diaminohexane ($pK_1=11$; $pK_2=10$), 1,3-diaminopentane (Dytek EP) ($pK_1=10.5$; $pK_2=8.9$), 2-methyl 1,5-diaminopentane (Dytek A) ($pK_1=11.2$; $pK_2=10.0$). Other preferred materials are the primary/primary diamines having alkylene spacers ranging from C_4 - C_8 . In general, primary diamines are preferred over secondary and tertiary diamines.

The following are non-limiting examples of diamines suitable for use in the present invention.


1-N,N-dimethylamino-3-aminopropane having the formula:




1,6-diaminohexane having the formula:




1,3-diaminopropane having the formula:




2-methyl-1,5-diaminopentane having the formula:




1,3-diaminopentane, available under the tradename Dytek EP, having the formula:




1,3-diaminobutane having the formula:




Jeffamine EDR 148, a diamine having an alkyleneoxy backbone, having the formula:



3-methyl-3-aminoethyl-5-dimethyl-1-aminocyclohexane (isophorone diamine) having the formula:



1,3-bis(methylamino)cyclohexane having the formula:



ADJUNCT INGREDIENTS

Builder - The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylene-diamine tetraacetate, metal ion sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylene-phosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also be used herein.

Suitable polycarboxylates builders for use herein include citric acid, preferably in the form of a water-soluble salt, derivatives of succinic acid of the formula $R-CH(COOH)CH_2(COOH)$ wherein R is C10-20 alkyl or alkenyl, preferably C12-16, or wherein R can be substituted with hydroxyl, sulfo sulfoxyl or sulfone substituents. Specific examples include lauryl succinate, myristyl succinate, palmityl succinate 2-dodecenylsuccinate, 2-tetradecenyl succinate. Succinate builders are preferably used in the form of their water-soluble salts, including sodium, potassium, ammonium and alkanolammonium salts.

Other suitable polycarboxylates are oxodisuccinates and mixtures of tartrate monosuccinic and tartrate disuccinic acid such as described in US 4,663,071.

Especially for the liquid execution herein, suitable fatty acid builders for use herein are saturated or unsaturated C10-18 fatty acids, as well as the corresponding soaps. Preferred saturated species have from 12 to 16 carbon atoms in the alkyl chain. The preferred unsaturated fatty acid is oleic acid. Other preferred builder system for liquid compositions is based on dodecenyl succinic acid and citric acid.

Detergency builder salts are normally included in amounts of from 3% to 50% by weight of the composition preferably from 5% to 30% and most usually from 5% to 25% by weight.

OPTIONAL DETERGENT INGREDIENTS:

Enzymes - Detergent compositions of the present invention may further comprise one or more enzymes which provide cleaning performance benefits. Said enzymes include enzymes selected from cellulases, hemicellulases, peroxidases, proteases, glucosylases, amylases, lipases, cutinases, pectinases, xylanases, reductases, oxidases, phenoloxidases, lipoxigenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases or mixtures thereof. A preferred combination is a detergent composition having a cocktail of conventional applicable enzymes like protease, amylase, lipase, cutinase and/or cellulase. Enzymes when present in the compositions, at from about 0.0001% to about 5% of active enzyme by weight of the detergent composition.

Proteolytic Enzyme - The proteolytic enzyme can be of animal, vegetable or microorganism (preferred) origin. The proteases for use in the detergent compositions herein include (but are not limited to) trypsin, subtilisin, chymotrypsin and elastase-type proteases. Preferred for use herein are subtilisin-type proteolytic enzymes. Particularly preferred is bacterial serine proteolytic enzyme obtained from Bacillus subtilis and/or Bacillus licheniformis.

Suitable proteolytic enzymes include Novo Industri A/S Alcalase[®] (preferred), Esperase[®], Savinase[®] (Copenhagen, Denmark), Gist-brocades' Maxatase[®], Maxacal[®] and Maxapem 15[®] (protein engineered Maxacal[®]) (Delft, Netherlands), and subtilisin BPN and BPN' (preferred), which are commercially available. Preferred proteolytic enzymes are also modified bacterial serine proteases, such as those made by Genencor International, Inc. (San Francisco, California) which are described in European Patent 251,446B, granted December 28, 1994 (particularly pages 17, 24 and 98) and which are also called herein "Protease B". U.S. Patent 5,030,378, Venegas, issued July 9, 1991, refers to a modified bacterial serine proteolytic enzyme (Genencor International) which is called "Protease A" herein (same as BPN'). In particular see columns 2 and 3 of U.S. Patent 5,030,378 for a complete description, including amino sequence, of Protease A and its variants. Other proteases are sold under the tradenames: Primase, Durazym, Opticlean and Optimase. Preferred proteolytic enzymes, then, are selected from the group consisting of Alcalase[®] (Novo Industri A/S), BPN', Protease A and Protease B (Genencor), and mixtures thereof. Protease B is most preferred.

Of particular interest for use herein are the proteases described in U.S. Patent No. 5,470,733.

Also proteases described in our co-pending application USSN 08/136,797 can be included in the detergent composition of the invention.

Another preferred protease, referred to as "Protease D" is a carbonyl hydrolase variant having an amino acid sequence not found in nature, which is derived from a precursor carbonyl hydrolase by substituting a different amino acid for a plurality of amino acid residues at a position in said carbonyl hydrolase equivalent to position +76, preferably also in combination with one or more amino acid residue positions equivalent to those selected from the group consisting of +99, +101, +103, +104, +107, +123, +27, +105, +109, +126, +128, +135, +156, +166, +195, +197, +204, +206, +210, +216, +217, +218, +222, +260, +265, and/or +274 according to the numbering of *Bacillus amyloliquefaciens* subtilisin, as described in WO 95/10615 published April 20, 1995 by Genencor International (A. Baeck et al. entitled "Protease-Containing Cleaning Compositions" having U.S. Serial No. 08/322,676, filed October 13, 1994).

Useful proteases are also described in PCT publications: WO 95/30010 published November 9, 1995 by The Procter & Gamble Company; WO 95/30011 published November 9, 1995 by The Procter & Gamble Company; WO 95/29979 published November 9, 1995 by The Procter & Gamble Company.

Protease enzyme may be incorporated into the compositions in accordance with the invention at a level of from 0.0001% to 2% active enzyme by weight of the composition.

Amylase - Amylases (α and/or β) can be included for removal of carbohydrate-based stains. Suitable amylases are Termamyl® (Novo Nordisk), Fungamyl® and BAN® (Novo Nordisk). The enzymes may be of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin. Amylase enzymes are normally incorporated in the detergent composition at levels from 0.0001% to 2%, preferably from about 0.0001% to about 0.5%, more preferably from about 0.0005% to about 0.1%, even more preferably from about 0.001% to about 0.05% of active enzyme by weight of the detergent composition.

Amylase enzymes also include those described in WO95/26397 and in co-pending application by Novo Nordisk PCT/DK96/00056. Other specific amylase enzymes for use in the detergent compositions of the present invention therefore include : (a) α -amylases characterised by having a specific activity at least 25% higher than the specific activity of Termamyl® at a temperature range of 25°C to 55°C and at a pH value in the range of 8 to 10, measured by the Phadebas® α -amylase activity assay. Such Phadebas® α -amylase activity assay is described at pages 9-10, WO95/26397.

(b) α -amylases according (a) comprising the amino sequence shown in the SEQ ID listings in the above cited reference. or an α -amylase being at least 80% homologous with the amino acid sequence shown in the SEQ ID listing.

(c) α -amylases according (a) obtained from an alkalophilic *Bacillus* species, comprising the following amino sequence in the N-terminal : His-His-Asn-Gly-Thr-Asn-Gly-Thr-Met-Met-Gln-Tyr-Phe-Glu-Trp-Tyr-Leu-Pro-Asn-Asp.

A polypeptide is considered to be X% homologous to the parent amylase if a comparison of the respective amino acid sequences, performed via algorithms, such as the one described by Lipman and Pearson in Science 227, 1985, p. 1435, reveals an identity of X%

(d) α -amylases according (a-c) wherein the α -amylase is obtainable from an alkalophilic *Bacillus* species; and in particular, from any of the strains NCIB 12289, NCIB 12512, NCIB 12513 and DSM 935.

In the context of the present invention, the term "obtainable from" is intended not only to indicate an amylase produced by a *Bacillus* strain but also an amylase encoded by a DNA sequence isolated from such a *Bacillus* strain and produced in an host organism transformed with said DNA sequence.

(e) α -amylase showing positive immunological cross-reactivity with antibodies raised against an α -amylase having an amino acid sequence corresponding respectively to those α -amylases in (a-d).

(f) Variants of the following parent α -amylases which (i) have one of the amino acid sequences shown in corresponding respectively to those α -amylases in (a-e), or (ii) displays at least 80% homology with one or more of said amino acid sequences, and/or displays immunological cross-reactivity with an antibody raised against an α -amylase having one of said amino acid sequences, and/or is encoded by a DNA sequence which hybridizes with the same probe as a DNA sequence encoding an α -amylase having one of said amino acid sequence; in which variants :

1. at least one amino acid residue of said parent α -amylase has been deleted; and/or
2. at least one amino acid residue of said parent α -amylase has been replaced by a different amino acid residue; and/or
3. at least one amino acid residue has been inserted relative to said parent α -amylase;

said variant having an α -amylase activity and exhibiting at least one of the following properties relative to said parent α -amylase : increased thermostability, increased stability towards oxidation, reduced Ca ion

dependency, increased stability and/or α -amylolytic activity at neutral to relatively high pH values, increased α -amylolytic activity at relatively high temperature and increase or decrease of the isoelectric point (pI) so as to better match the pI value for α -amylase variant to the pH of the medium.

Said variants are described in the patent application PCT/DK96/00056.

Other amylases suitable herein include, for example, α -amylases described in GB 1,296,839 to Novo; RAPIDASE®, International Bio-Synthetics, Inc. and TERMAMYL®, Novo. FUNGAMYL® from Novo is especially useful. Engineering of enzymes for improved stability, e.g., oxidative stability, is known. See, for example J. Biological Chem., Vol. 260, No. 11, June 1985, pp. 6518-6521. Certain preferred embodiments of the present compositions can make use of amylases having improved stability in detergents such as automatic dishwashing types, especially improved oxidative stability as measured against a reference-point of TERMAMYL® in commercial use in 1993. These preferred amylases herein share the characteristic of being "stability-enhanced" amylases, characterized, at a minimum, by a measurable improvement in one or more of: oxidative stability, e.g., to hydrogen peroxide/tetraacetythylenediamine in buffered solution at pH 9-10; thermal stability, e.g., at common wash temperatures such as about 60°C; or alkaline stability, e.g., at a pH from about 8 to about 11, measured versus the above-identified reference-point amylase. Stability can be measured using any of the art-disclosed technical tests. See, for example, references disclosed in WO 9402597. Stability-enhanced amylases can be obtained from Novo or from Genencor International. One class of highly preferred amylases herein have the commonality of being derived using site-directed mutagenesis from one or more of the Bacillus amylases, especially the Bacillus α -amylases, regardless of whether one, two or multiple amylase strains are the immediate precursors. Oxidative stability-enhanced amylases vs. the above-identified reference amylase are preferred for use, especially in bleaching, more preferably oxygen bleaching, as distinct from chlorine bleaching, detergent compositions herein. Such preferred amylases include (a) an amylase according to the hereinbefore incorporated WO 9402597, Novo, Feb. 3, 1994, as further illustrated by a mutant in which substitution is made, using alanine or threonine, preferably threonine, of the methionine residue located in position 197 of the B. licheniformis alpha-amylase, known as TERMAMYL®, or the homologous position variation of a similar parent amylase, such as B. amyloliquefaciens, B. subtilis, or B. stearothermophilus; (b) stability-enhanced amylases as described by Genencor International in a paper entitled "Oxidatively Resistant alpha-Amylases" presented at the 207th American Chemical Society National Meeting, March 13-17 1994, by C.

Mitchinson. Therein it was noted that bleaches in automatic dishwashing detergents inactivate alpha-amylases but that improved oxidative stability amylases have been made by Genencor from *B. licheniformis* NCIB8061. Methionine (Met) was identified as the most likely residue to be modified. Met was substituted, one at a time, in positions 8, 15, 197, 256, 304, 366 and 438 leading to specific mutants, particularly important being M197L and M197T with the M197T variant being the most stable expressed variant. Stability was measured in CASCADE® and SUNLIGHT®; (c) particularly preferred amylases herein include amylase variants having additional modification in the immediate parent as described in WO 9510603 A and are available from the assignee, Novo, as DURAMYL®. Other particularly preferred oxidative stability enhanced amylase include those described in WO 9418314 to Genencor International and WO 9402597 to Novo. Any other oxidative stability-enhanced amylase can be used, for example as derived by site-directed mutagenesis from known chimeric, hybrid or simple mutant parent forms of available amylases. Other preferred enzyme modifications are accessible. See WO 9509909 A to Novo.

Various carbohydrase enzymes which impart antimicrobial activity may also be included in the present invention. Such enzymes include endoglycosidase, Type II endoglycosidase and glucosidase as disclosed in U.S. Patent Nos. 5,041,236, 5,395,541, 5,238,843 and 5,356,803 the disclosures of which are herein incorporated by reference. Of course, other enzymes having antimicrobial activity may be employed as well including peroxidases, oxidases and various other enzymes.

It is also possible to include an enzyme stabilization system into the compositions of the present invention when any enzyme is present in the composition.

Perfumes - Perfumes and perfumery ingredients useful in the present compositions and processes comprise a wide variety of natural and synthetic chemical ingredients, including, but not limited to, aldehydes, ketones, esters, and the like. Also included are various natural extracts and essences which can comprise complex mixtures of ingredients, such as orange oil, lemon oil, rose extract, lavender, musk, patchouli, balsamic essence, sandalwood oil, pine oil, cedar, and the like. Finished perfumes can comprise extremely complex mixtures of such ingredients. Finished perfumes typically comprise from about 0.01% to about 2%, by weight, of the detergent compositions herein, and individual perfumery ingredients can comprise from about 0.0001% to about 90% of a finished perfume composition.

Non-limiting examples of perfume ingredients useful herein include: 7-acetyl-1,2,3,4,5,6,7,8-octahydro-1,1,6,7-tetramethyl naphthalene; ionone methyl; ionone gamma methyl; methyl cedrylone; methyl dihydrojasmonate; methyl 1,6,10-trimethyl-

2,5,9-cyclododecatrien-1-yl ketone; 7-acetyl-1,1,3,4,4,6-hexamethyl tetralin; 4-acetyl-6-tert-butyl-1,1-dimethyl indane; para-hydroxy-phenyl-butanone; benzophenone; methyl beta-naphthyl ketone; 6-acetyl-1,1,2,3,3,5-hexamethyl indane; 5-acetyl-3-isopropyl-1,1,2,6-tetramethyl indane; 1-dodecanal, 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde; 7-hydroxy-3,7-dimethyl ocatanal; 10-undecen-1-al; iso-hexenyl cyclohexyl carboxaldehyde; formyl tricyclodecane; condensation products of hydroxycitronellal and methyl anthranilate, condensation products of hydroxycitronellal and indol, condensation products of phenyl acetaldehyde and indol; 2-methyl-3-(para-tert-butylphenyl)-propionaldehyde; ethyl vanillin; heliotropin; hexyl cinnamic aldehyde; amyl cinnamic aldehyde; 2-methyl-2-(para-iso-propylphenyl)-propionaldehyde; coumarin; decalactone gamma; cyclopentadecanolide; 16-hydroxy-9-hexadecenoic acid lactone; 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta-gamma-2-benzopyrane; beta-naphthol methyl ether; ambroxane; dodecahydro-3a,6,6,9a-tetramethylnaphtho[2,1b]furan; cedrol, 5-(2,2,3-trimethylcyclopent-3-enyl)-3-methylpentan-2-ol; 2-ethyl-4-(2,2,3-trimethyl-3-cyclopenten-1-yl)-2-buten-1-ol; caryophyllene alcohol; tricyclodecenyl propionate; tricyclodecenyl acetate; benzyl salicylate; cedryl acetate; and para-(tert-butyl) cyclohexyl acetate.

Particularly preferred perfume materials are those that provide the largest odor improvements in finished product compositions containing cellulases. These perfumes include but are not limited to: hexyl cinnamic aldehyde; 2-methyl-3-(para-tert-butylphenyl)-propionaldehyde; 7-acetyl-1,2,3,4,5,6,7,8-octahydro-1,1,6,7-tetramethyl naphthalene; benzyl salicylate; 7-acetyl-1,1,3,4,4,6-hexamethyl tetralin; para-tert-butyl cyclohexyl acetate; methyl dihydro jasmonate; beta-naphthol methyl ether; methyl beta-naphthyl ketone; 2-methyl-2-(para-iso-propylphenyl)-propionaldehyde; 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-cyclopenta-gamma-2-benzopyrane; dodecahydro-3a,6,6,9a-tetramethylnaphtho[2,1b]furan; anisaldehyde; coumarin; cedrol; vanillin; cyclopentadecanolide; tricyclodecenyl acetate; and tricyclodecenyl propionate.

Other perfume materials include essential oils, resinoids, and resins from a variety of sources including, but not limited to: Peru balsam, Olibanum resinoid, styrax, labdanum resin, nutmeg, cassia oil, benzoin resin, coriander and lavandin. Still other perfume chemicals include phenyl ethyl alcohol, terpineol, linalool, linalyl acetate, geraniol, nerol, 2-(1,1-dimethylethyl)-cyclohexanol acetate, benzyl acetate, and eugenol. Carriers such as diethylphthalate can be used in the finished perfume compositions.

Chelating Agents - The detergent compositions herein may also optionally contain one or more iron and/or manganese chelating agents. Such chelating agents can be selected from the group consisting of amino carboxylates, amino phosphonates,

polyfunctionally-substituted aromatic chelating agents and mixtures therein, all as hereinafter defined. Without intending to be bound by theory, it is believed that the benefit of these materials is due in part to their exceptional ability to remove iron and manganese ions from washing solutions by formation of soluble chelates.

Amino carboxylates useful as optional chelating agents include ethylenediaminetetraacetates, N-hydroxyethylethylenediaminetriacetates, nitrilo-triacetates, ethylenediamine tetrapropionates, triethylenetetraaminehexacetates, diethylenetriaminepentaacetates, and ethanoldi-glycines, alkali metal, ammonium, and substituted ammonium salts therein and mixtures therein.

Amino phosphonates are also suitable for use as chelating agents in the compositions of the invention when at least low levels of total phosphorus are permitted in detergent compositions, and include ethylenediaminetetrakis (methylenephosphonates) as DEQUEST. Preferred, these amino phosphonates do not contain alkyl or alkenyl groups with more than about 6 carbon atoms.

Polyfunctionally-substituted aromatic chelating agents are also useful in the compositions herein. See U.S. Patent 3,812,044, issued May 21, 1974, to Connor et al. Preferred compounds of this type in acid form are dihydroxydisulfobenzenes such as 1,2-dihydroxy-3,5-disulfobenzene.

A preferred biodegradable chelator for use herein is ethylenediamine disuccinate ("EDDS"), especially the [S,S] isomer as described in U.S. Patent 4,704,233, November 3, 1987, to Hartman and Perkins.

The compositions herein may also contain water-soluble methyl glycine diacetic acid (MGDA) salts (or acid form) as a chelant or co-builder. Similarly, the so called "weak" builders such as citrate can also be used as chelating agents.

If utilized, these chelating agents will generally comprise from about 0.1% to about 15% by weight of the detergent compositions herein. More preferably, if utilized, the chelating agents will comprise from about 0.1% to about 3.0% by weight of such compositions.

Composition pH

Dishwashing compositions of the invention will be subjected to acidic stresses created by food soils when put to use, i.e., diluted and applied to soiled dishes. If a composition with a pH greater than 7 is to be more effective, it preferably should contain a buffering agent capable of providing a generally more alkaline pH in the composition and in dilute solutions, i.e., about 0.1% to 0.4% by weight aqueous solution, of the composition. The pKa value of this buffering agent should be about 0.5 to 1.0 pH units below the desired pH value of the composition (determined as described above).

Preferably, the pKa of the buffering agent should be from about 7 to about 10. Under these conditions the buffering agent most effectively controls the pH while using the least amount thereof.

The buffering agent may be an active detergent in its own right, or it may be a low molecular weight, organic or inorganic material that is used in this composition solely for maintaining an alkaline pH. Preferred buffering agents for compositions of this invention are nitrogen-containing materials. Some examples are amino acids such as lysine or lower alcohol amines like mono-, di-, and tri-ethanolamine. Other preferred nitrogen-containing buffering agents are Tri(hydroxymethyl)amino methane ($(\text{HOCH}_2)_3\text{CNH}_2$ (TRIS), 2-amino-2-ethyl-1,3-propanediol, 2-amino-2-methyl-propanol, 2-amino-2-methyl-1,3-propanol, disodium glutamate, N-methyl diethanolamide, 1,3-diamino-propanol, N,N'-tetra-methyl-1,3-diamino-2-propanol, N,N-bis(2-hydroxyethyl)glycine (bicine) and N-tris (hydroxymethyl)methyl glycine (tricine). Mixtures of any of the above are also acceptable. Useful inorganic buffers/alkalinity sources include the alkali metal carbonates and alkali metal phosphates, e.g., sodium carbonate, sodium polyphosphate. For additional buffers see McCutcheon's EMULSIFIERS AND DETERGENTS, North American Edition, 1997, McCutcheon Division, MC Publishing Company Kirk and WO 95/07971 both of which are incorporated herein by reference.

The buffering agent, if used, is present in the compositions of the invention herein at a level of from about 0.1% to 15%, preferably from about 1% to 10%, most preferably from about 2% to 8%, by weight of the composition.

Calcium and/or Magnesium Ions

The presence of calcium and/or magnesium (divalent) ions improves the cleaning of greasy soils for various compositions, i.e., compositions containing alkyl ethoxy sulfates and/or polyhydroxy fatty acid amides. This is especially true when the compositions are used in softened water that contains few divalent ions. It is believed that calcium and/or magnesium ions increase the packing of the surfactants at the oil/water interface, thereby reducing interfacial tension and improving grease cleaning.

Compositions of the invention herein containing magnesium and/or calcium ions exhibit good grease removal, manifest mildness to the skin, and provide good storage stability. These ions can be present in the compositions herein at an active level of from about 0.1% to 4%, preferably from about 0.3% to 3.5%, more preferably from about 0.5% to 1%, by weight.

Preferably, the magnesium or calcium ions are added as a hydroxide, chloride, acetate, formate, oxide or nitrate salt to the compositions of the present invention. Calcium ions may also be added as salts of the hydrotrope.

The amount of calcium or magnesium ions present in compositions of the invention will be dependent upon the amount of total surfactant present therein. When calcium ions are present in the compositions of this invention, the molar ratio of calcium ions to total anionic surfactant should be from about 0.25:1 to about 2:1.

Formulating such divalent ion-containing compositions in alkaline pH matrices may be difficult due to the incompatibility of the divalent ions, particularly magnesium, with hydroxide ions. When both divalent ions and alkaline pH are combined with the surfactant mixture of this invention, grease cleaning is achieved that is superior to that obtained by either alkaline pH or divalent ions alone. Yet, during storage, the stability of these compositions becomes poor due to the formation of hydroxide precipitates. Therefore, chelating agents discussed hereinbefore may also be necessary.

Other Ingredients - The detergent compositions will further preferably comprise one or more deterative adjuncts selected from the following: soil release polymers, polymeric dispersants, polysaccharides, abrasives, bactericides, tarnish inhibitors, builders, enzymes, opacifiers, dyes, buffers, antifungal or mildew control agents, insect repellents, perfumes, hydrotropes, thickeners, processing aids, suds boosters, brighteners, anti-corrosive aids, stabilizers antioxidants and chelants. A wide variety of other ingredients useful in detergent compositions can be included in the compositions herein, including other active ingredients, carriers, hydrotropes, antioxidants, processing aids, dyes or pigments, solvents for liquid formulations, solid fillers for bar compositions, etc. If high sudsing is desired, suds boosters such as the C₁₀-C₁₆ alkanolamides can be incorporated into the compositions, typically at 1%-10% levels. The C₁₀-C₁₄ monoethanol and diethanol amides illustrate a typical class of such suds boosters. Use of such suds boosters with high sudsing adjunct surfactants such as the amine oxides, betaines and sultaines noted above is also advantageous.

An antioxidant can be optionally added to the detergent compositions of the present invention. They can be any conventional antioxidant used in detergent compositions, such as 2,6-di-tert-butyl-4-methylphenol (BHT), carbamate, ascorbate, thiosulfate, monoethanolamine(MEA), diethanolamine, triethanolamine, etc. It is preferred that the antioxidant, when present, be present in the composition from about 0.001% to about 5% by weight.

Various deterative ingredients employed in the present compositions optionally can be further stabilized by absorbing said ingredients onto a porous hydrophobic

substrate, then coating said substrate with a hydrophobic coating. Preferably, the deterative ingredient is admixed with a surfactant before being absorbed into the porous substrate. In use, the deterative ingredient is released from the substrate into the aqueous washing liquor, where it performs its intended deterative function.

To illustrate this technique in more detail, a porous hydrophobic silica (trademark SIPERNAT D10, DeGussa) is admixed with a proteolytic enzyme solution containing 3%-5% of C₁₃₋₁₅ ethoxylated alcohol (EO 7) nonionic surfactant. Typically, the enzyme/surfactant solution is 2.5 X the weight of silica. The resulting powder is dispersed with stirring in silicone oil (various silicone oil viscosities in the range of 500-12,500 can be used). The resulting silicone oil dispersion is emulsified or otherwise added to the final detergent matrix. By this means, ingredients such as the aforementioned enzymes, bleaches, bleach activators, bleach catalysts, photoactivators, dyes, fluorescers, fabric conditioners and hydrolyzable surfactants can be "protected" for use in detergents, including liquid laundry detergent compositions.

Further, these hand dishwashing detergent embodiments preferably further comprises a hydrotrope. Suitable hydrotropes include sodium, potassium, ammonium or water-soluble substituted ammonium salts of toluene sulfonic acid, naphthalene sulfonic acid, cumene sulfonic acid, xylene sulfonic acid.

The detergent compositions of this invention can be in any form, including granular, paste, gel or liquid. Highly preferred embodiments are in liquid or gel form. Liquid detergent compositions can contain water and other solvents as carriers. Low molecular weight primary or secondary alcohols exemplified by methanol, ethanol, propanol, and isopropanol are suitable. Monohydric alcohols are preferred for solubilizing surfactant, but polyols such as those containing from 2 to about 6 carbon atoms and from 2 to about 6 hydroxy groups (e.g., 1,3-propanediol, ethylene glycol, glycerine, and 1,2-propanediol) can also be used. The compositions may contain from 5% to 90%, typically 10% to 50% of such carriers.

An example of the procedure for making granules of the detergent compositions herein is as follows: - Linear alkylbenzenesulfonate, citric acid, sodium silicate, sodium sulfate perfume, diamine and water are added to, heated and mixed via a crutcher. The resulting slurry is spray dried into a granular form.

An example of the procedure for making liquid detergent compositions herein is as follows: - To the free water and citrate are added and dissolved. To this solution amine oxide, betaine, ethanol, hydrotrope and nonionic surfactant are added. If free water isn't available, the citrate are added to the above mix then stirred until dissolved. At this point, an acid is added to neutralize the formulation. It is preferred that the acid

be chosen from organic acids such as maleic and citric, however, inorganic mineral acids may be employed as well. In preferred embodiments these acids are added to the formulation followed by diamine addition. AExS is added last.

Non-Aqueous Liquid Detergents

The manufacture of liquid detergent compositions which comprise a non-aqueous carrier medium can be prepared according to the disclosures of U.S. Patents 4,753,570; 4,767,558; 4,772,413; 4,889,652; 4,892,673; GB-A-2,158,838; GB-A-2,195,125; GB-A-2,195,649; U.S. 4,988,462; U.S. 5,266,233; EP-A-225,654 (6/16/87); EP-A-510,762 (10/28/92); EP-A-540,089 (5/5/93); EP-A-540,090 (5/5/93); U.S. 4,615,820; EP-A-565,017 (10/13/93); EP-A-030,096 (6/10/81), incorporated herein by reference. Such compositions can contain various particulate deterative ingredients stably suspended therein. Such non-aqueous compositions thus comprise a LIQUID PHASE and, optionally but preferably, a SOLID PHASE, all as described in more detail hereinafter and in the cited references.

The compositions of this invention can be used to form aqueous washing solutions for use hand dishwashing. Generally, an effective amount of such compositions is added to water to form such aqueous cleaning or soaking solutions. The aqueous solution so formed is then contacted with the dishware, tableware, and cooking utensils.

An effective amount of the detergent compositions herein added to water to form aqueous cleaning solutions can comprise amounts sufficient to form from about 500 to 20,000 ppm of composition in aqueous solution. More preferably, from about 800 to 5,000 ppm of the detergent compositions herein will be provided in aqueous cleaning liquor.

METHOD OF USE

The present invention also relates to a method for providing increased suds volume and increased suds retention while hand washing dishware or cookware articles in need of cleaning, comprising the step of contacting said articles with an aqueous solution of a detergent composition suitable for use in hand dishwashing, said composition comprising:

- a) an effective amount of a polymeric suds stabilizer as hereinbefore defined;
 - b) an effective amount of a deterative surfactant; and
 - c) the balance carriers and other adjunct ingredients;
- provided the pH of a 10% aqueous solution of said composition is from about 4 to about 12.

The present invention also relates to a means for preventing the redeposition of grease, oils, and dirt, especially grease, from the hand washing solution onto dishware. This method comprises contacting an aqueous solution of the compositions of the present invention with soiled dishware and washing said dishware with said aqueous solution.

An effective amount of the detergent compositions herein added to water to form aqueous cleaning solutions according to the method of the present invention comprises amounts sufficient to form from about 500 to 20,000 ppm of composition in aqueous solution. More preferably, from about 800 to 2,500 ppm of the detergent compositions herein will be provided in aqueous cleaning liquor.

The liquid detergent compositions of the present invention are effective for preventing the redeposition of grease from the wash solution back onto the dishware during washing. One measure of effectiveness of the compositions of the present invention involves redeposition tests. The following test and others of similar nature are used to evaluate the suitability of the formulas described herein.

A polyethylene 2 L graduated cylinder is filled to the 1 L graduation mark with an aqueous (water = 7 grain) solution comprising from about 500 to about 20,000 ppm of a liquid detergent composition according to the present invention. A synthetic greasy soil composition is then added to the cylinder and the solution is agitated. After a period of time the solution is decanted from the graduated cylinder and the interior walls of the graduated cylinder are rinsed with a suitable solvent or combination of solvents to recover any re-deposited greasy soil. The solvent is removed and the weight of greasy soil which remains in solution is determined by subtracting the amount of soil recovered from the amount initially added to the aqueous solution.

Other re-deposition test include immersion of tableware, flatware, and the like and recovering any re-deposited soil.

The above test can be further modified to determine the increased amount of suds volume and suds duration. The solution is first agitated then subsequently challenged with portions of greasy soil with agitation between each subsequent soil addition. The suds volume can be easily determined by using the vacant volume of the 2 L cylinder as a guide.

The following are non-limiting examples of liquid detergent compositions comprising the polymeric suds extenders according to the present invention.

EXAMPLE 1

Reaction of (1-octene/maleic anhydride) copolymer with 1 equivalent of DMAPA

Poly(maleic anhydride-alt-1-octene) (15.00 g) and tetrahydrofuran (200 ml, anhydrous) are placed into a 250 ml three-necked round-bottom flask, fitted with a heating mantle, magnetic stirrer, dropping funnel, internal thermometer and argon inlet. 3-Dimethylaminopropylamine (7.65 g, 74.87 mmol) is added dropwise over 15 minutes, with an exotherm to 30°C and heavy precipitation. The mixture is stirred for 4 hours at 55°C. The mixture is poured into 3:1 ethyl ether:hexanes to precipitate the product which is dried under vacuum to yield a white powder. NMR is consistent with the desired compound.

EXAMPLE 2

Reaction of (1-hexene/maleic anhydride) copolymer with 1 equivalent of DMAPA

Poly(maleic anhydride-alt-1-hexene) (15.00 g) and pyridine (150 ml, anhydrous) are placed into a 250 ml three-necked round-bottom flask, fitted with a heating mantle, magnetic stirrer, dropping funnel, internal thermometer and argon inlet. There is a slight exotherm and the mixture is dark. 3-Dimethylaminopropylamine (9.25 g, 90.53 mmol) is added dropwise over 15 minutes, with an exotherm to 45°C. The mixture is stirred for 4 hours at 80°C. The mixture is concentrated by rotary evaporation, dissolved into water and lyophilized to yield a yellow powder. NMR is consistent with the desired compound.

EXAMPLE 3

Preparation of Poly(DMAM-co-AA) (2:1) Copolymer

2-(Dimethylamino)ethyl methacrylate (90.00 g, 572.4 mmol), acrylic acid (20.63g, 286.2 mmol), 2,2'-azobisisobutyronitrile (0.70 g, 4.3 mmol), 1,4-dioxane (345 ml) and 2-propanol (86 ml) are placed into a 1000 ml three-necked round-bottomed flask, fitted with a heating mantle, magnetic stirrer, internal thermometer and argon inlet. The mixture is sparged with nitrogen for 30 minutes to remove dissolved oxygen. The mixture is heated for 18 hours with stirring at 65°C. TLC (diethyl ether) indicates consumption of monomer. The mixture is concentrated under vacuum by rotary evaporation to remove the solvent. Water is added to make a 10% solution and the mixture is lyophilized and then pulverized in a blender to yield an off-white-peach powder. NMR is consistent with the desired compound.

The following are non-limiting examples of liquid detergent compositions comprising the polymeric suds extenders according to the present invention.

TABLE I

weight %

Ingredients	4	5	6
C ₁₂ -C ₁₅ Alkyl sulphate	--	28.0	25.0
C ₁₂ -C ₁₃ Alkyl (E _{0.6-3}) sulfate	30	--	--
C ₁₂ Amine oxide	5.0	3.0	7.0
C ₁₂ -C ₁₄ Betaine	3.0	--	1.0
C ₁₂ -C ₁₄ Polyhydroxy fatty acid amide	--	1.5	--
C ₁₀ Alcohol Ethoxylate E ₉ ¹	2.0	--	4.0
Diamine ²	0.5	--	7.0
Mg ²⁺ (as MgCl ₂)	0.25	--	--
Citrate (cit2K3)	0.25	--	--
Polymeric suds booster ³	1.25	2.6	0.9
Minors and water ⁴	balance	balance	balance
pH of a 10% aqueous solution	9	10	10

1. E₉ Ethoxylated Alcohols as sold by the Shell Oil Co.

2. 1,3-bis(methylamino)cyclohexane.

3. Copolymer of Example 3.

4. Includes perfumes, dyes, ethanol, etc.

TABLE II

Ingredients	weight %		
	7	8	9
C ₁₂ -C ₁₃ Alkyl (E _{0.6-3}) sulfate	--	15.0	10.0
Paraffin sulfonate	20.0	--	--
Na C ₁₂ -C ₁₃ linear alkylbenzene sulfonate	5.0	15.0	12.0
C ₁₂ -C ₁₄ Betaine	3.0	1.0	--
C ₁₂ -C ₁₄ Polyhydroxy fatty acid amide	3.0	--	1.0
C ₁₀ Alcohol Ethoxylate E ₉ ¹	--	--	20.0
Diamine ²	1.0	-	7.0
DTPA ³	--	0.2	--
Mg ²⁺ (as MgCl ₂)	1.0	--	--
Ca ²⁺ (as Ca(citrate) ₂)	--	0.5	--
Protease ⁴	0.01	--	0.05
Amylase ⁵	--	0.05	0.05
Hydrotrope ⁶	2.0	1.5	3.0
Polymeric suds booster ⁷	0.1	3.0	0.5

Minors and water ⁸	balance	balance	balance
pH of a 10% aqueous solution	9.3	8.5	11

1. E₉ Ethoxylated Alcohols as sold by the Shell Oil Co.
2. 1,3-diaminopentane sold as Dytek EP.
3. Diethylenetriaminepentaacetate.
4. Suitable protease enzymes include Savinase[®]; Maxatase[®]; Maxacal[®]; Maxapem 15[®]; subtilisin BPN and BPN'; Protease B; Protease A; Protease D; Primase[®]; Durazym[®]; Opticlean[®]; and Optimase[®]; and Alcalase[®].
5. Suitable amylase enzymes include Termamyl[®], Fungamyl[®]; Duramyl[®]; BAN[®], and the amylases as described in WO95/26397 and in co-pending application by Novo Nordisk PCT/DK/96/00056.
6. Suitable hydrotropes include sodium, potassium, ammonium or water-soluble substituted ammonium salts of toluene sulfonic acid, naphthalene sulfonic acid, cumene sulfonic acid, xylene sulfonic acid.
7. LX1279 available from Baker Petrolite.
8. Includes perfumes, dyes, ethanol, etc.

TABLE III

Ingredients	weight %			
	10	11	12	13
C ₁₂ -C ₁₅ Alkyl (E ₁) sulfate	--	30.0	--	--
C ₁₂ -C ₁₅ Alkyl (E _{1,4}) sulfate	30.0	--	27.0	--
C ₁₂ -C ₁₅ Alkyl (E _{2,2}) sulfate	--	--	--	15
C ₁₂ Amine oxide	5.0	5.0	5.0	3.0
C ₁₂ -C ₁₄ Betaine	3.0	3.0	--	--
C ₁₀ Alcohol Ethoxylate E ₉ ¹	2.0	2.0	2.0	2.0
Diamine ²	1.0	2.0	4.0	2.0
Mg ²⁺ (as MgCl ₂)	0.25	0.25	--	--
Ca ²⁺ (as Ca(citrate) ₂)	--	0.4	--	--
Polymeric suds booster ³	0.2	1.0	0.75	5.0
Minors and water ⁴	balance	balance	balance	balance
pH of a 10% aqueous solution	7.4	7.6	7.4	7.8

1. E₉ Ethoxylated Alcohols as sold by the Shell Oil Co.
2. 1,3-diaminopentane sold as Dytek EP.
3. Copolymer of Example 2.

4. Includes perfumes, dyes, ethanol, etc.

TABLE IV

Ingredients	weight %		
	14	15	16
C ₁₂ -C ₁₃ Alkyl (E _{0.6-3}) sulfate	--	15.0	10.0
Paraffin sulfonate	20.0	--	--
Na C ₁₂ -C ₁₃ linear alkylbenzene sulfonate	5.0	15.0	12.0
C ₁₂ -C ₁₄ Betaine	3.0	1.0	--
C ₁₂ -C ₁₄ Polyhydroxy fatty acid amide	3.0	--	1.0
C ₁₀ Alcohol Ethoxylate E _g ¹	--	--	20.0
Diamine ²	1.0	--	7.0
Mg ²⁺ (as MgCl ₂)	1.0	--	--
Ca ²⁺ (as Ca(citrate) ₂)	--	0.5	--
Protease ³	0.1	--	--
Amylase ⁴	--	0.02	--
Lipase ⁵	--	--	0.025
DTPA ⁶	--	0.3	--
Citrate (cit2K3)	0.65	--	--
Polymeric suds booster ⁷	1.5	2.2	3.0
Minors and water ⁸	balance	balance	balance
pH of a 10% aqueous solution	9.3	8.5	11

1. E_g Ethoxylated Alcohols as sold by the Shell Oil Co.
2. 1,3-diaminopentane sold as Dytek EP.
3. Suitable protease enzymes include Savinase[®]; Maxatase[®]; Maxacal[®]; Maxapem 15[®]; subtilisin BPN and BPN'; Protease B; Protease A; Protease D; Primase[®]; Durazym[®]; Opticlean[®]; and Optimase[®]; and Alcalase[®].
4. Suitable amylase enzymes include Termamyl[®]; Fungamyl[®]; Duramyl[®]; BAN[®], and the amylases as described in WO95/26397 and in co-pending application by Novo Nordisk PCT/DK/96/00056.
5. Suitable lipase enzymes include Amano-P; M1 Lipase[®]; Lipomax[®]; Lipolase[®]; D96L - lipolytic enzyme variant of the native lipase derived from *Humicola lanuginosa* as described in US Patent Application Serial No. 08/341,826; and the *Humicola lanuginosa* strain DSM 4106
6. Diethylenetriaminepentaacetate.

7. Copolymer of Example 3.
8. Includes perfumes, dyes, ethanol, etc.

TABLE V

Ingredients	weight %		
	17	18	19
C ₁₂ -C ₁₃ Alkyl (E _{0.6-3}) sulfate	--	27.0	--
C ₁₂ -C ₁₄ Betaine	2.0	2.0	--
C ₁₄ Amine oxide	2.0	5.0	7.0
C ₁₂ -C ₁₄ Polyhydroxy fatty acid amide	2.0	--	--
C ₁₀ Alcohol Ethoxylate E ₉ ¹	1.0	--	2.0
Hydrotrope	--	--	5.0
Diamine ²	4.0	2.0	5.0
Ca ²⁺ (as Ca(citrate) ₂)	--	0.1	0.1
Protease ³	--	0.06	0.1
Amylase ⁴	0.005	--	0.05
Lipase ⁵	--	0.05	--
DTPA ⁶	--	0.1	0.1
Citrate (cit2K3)	0.3	--	--
Polymeric suds booster ⁷	0.5	0.8	0.3
Minors and water ⁸	balance	balance	balance
pH of a 10% aqueous solution	10	9	9.2

1. E₉ Ethoxylated Alcohols as sold by the Shell Oil Co.
2. 1,3-bis(methylamino)cyclohexane,
3. Suitable protease enzymes include Savinase[®]; Maxatase[®]; Maxacal[®]; Maxapem 15[®]; subtilisin BPN and BPN'; Protease B; Protease A; Protease D; Primase[®]; Durazym[®]; Opticlean[®]; and Optimase[®]; and Alcalase[®].
4. Suitable amylase enzymes include Termamyl[®], Fungamyl[®], Duramyl[®], BAN[®], and the amylases as described in WO95/26397 and in co-pending application by Novo Nordisk PCT/DK/96/00056.
5. Suitable lipase enzymes include Amano-P; M1 Lipase[®]; Lipomax[®]; Lipolase[®]; D96L - lipolytic enzyme variant of the native lipase derived from *Humicola lanuginosa* as described in US Patent Application Serial No. 08/341,826; and the *Humicola lanuginosa* strain DSM 4106
6. Diethylenetriaminepentaacetate.

7. Copolymer of Example 1.
8. Includes perfumes, dyes, ethanol, etc.


TABLE VI

Ingredients	weight %		
	20	21	22
C ₁₂ -C ₁₃ Alkyl (E _{1.4}) sulfate	33.29	24.0	—
C ₁₂ -C ₁₃ Alkyl (E _{0.6}) sulfate	--	--	26.26
C ₁₂ -C ₁₄ Polyhydroxy fatty acid amide	4.2	3.0	1.37
C ₁₄ Amine oxide	4.8	2.0	1.73
C ₁₁ Alcohol Ethoxylate E ₉ ¹	1.0	4.0	4.56
C ₁₂ -C ₁₄ Betaine	--	2.0	1.73
MgCl ₂	0.72	0.47	0.46
Calcium citrate	0.35	--	--
Polymeric suds booster ²	0.5	1.0	2.0
Minors and water ³	balance	balance	balance
pH of a 10% aqueous solution	7.4	7.8	7.8

1. E₉ Ethoxylated Alcohols as sold by the Shell Oil Co.
2. LX1279 available from Baker Petrolite.
3. Includes perfumes, dyes, ethanol, etc.


What is claimed is:

1. A detergent composition suitable for use in hand dishwashing, said composition comprising:
 - a) an effective amount of a zwitterionic polymeric suds stabilizer;
 - b) an effective amount of a deterative surfactant; and
 - c) the balance carriers and other adjunct ingredients;
 provided the pH of a 10% aqueous solution of said composition is from 4 to 12.
2. A composition according to Claim 1 wherein said zwitterionic polymeric suds stabilizer has the formula:




wherein R is C₁-C₁₂ linear alkylene, C₁-C₁₂ branched alkylene, and mixtures thereof; R¹ is a unit capable of having a negative charge at a pH of from 4 to 12; R² is a unit capable of having a positive charge at a pH of from 4 to 12; n has a value such that said zwitterionic polymers suds stabilizer has an average molecular weight of from 1,000 to 2,000,000 daltons; x is from 0 to 6; y is 0 or 1; and z is 0 or 1.

3. A composition according to any of Claims 1-2 wherein said zwitterionic polymeric suds stabilizer has an average molecular weight of from 5,000 to 1,000,000 daltons.
4. A composition according to any of Claims 2-3 wherein R¹ has the formula:




wherein L is a linking unit independently selected from the following:




and mixtures thereof; R' is independently hydrogen, C₁-C₄ alkyl, and mixtures thereof or R' and S can form a heterocycle of 4 to 7 carbon atoms, optionally containing other hetero atoms and optionally substituted; R³ is independently selected from -CO₂M, -SO₃M, -OSO₃M, -CH₂P(O)(OM)₂, -OP(O)(OM)₂, units having the formula:



wherein each R⁸, R⁹, and R¹⁰ is independently selected from the group consisting of hydrogen, -(CH₂)_mR¹¹, and mixtures thereof, wherein R¹¹ is -CO₂H, -SO₃M, -OSO₃M, -CH(CO₂H)CH₂CO₂H, -CH₂P(O)(OH)₂, -OP(O)(OH)₂, and mixtures thereof; provided that one R⁸, R⁹, or R¹⁰ is not a hydrogen atom; R² has the formula:



wherein L¹ is a linking unit independently selected from the following:




and mixtures thereof; wherein R' is independently hydrogen, C₁-C₄ alkyl, and mixtures thereof or alternatively R' and S can form a heterocycle of 4 to 7 carbon atoms, optionally containing other hetero atoms and optionally substituted; R⁴ is independently selected from amino, alkylamino carboxamide, 3-imidazolyl, 4-imidazolyl, 2-imidazolyl, 4-imidazolyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, piperazinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, guanidino, amidino, and mixtures


thereof; each S is independently selected from C₁-C₁₂ linear alkylene, C₁-C₁₂ branched alkylene, C₃-C₁₂ linear alkenylene, C₃-C₁₂ branched alkenylene, C₃-C₁₂ hydroxyalkylene, C₄-C₁₂ dihydroxyalkylene, C₆-C₁₀ arylene, C₈-C₁₂ dialkylarylene, $-(R^5O)_kR^5-$, $-(R^5O)_kR^6(OR^5)_k-$, $-CH_2CH(OR^7)CH_2-$, and mixtures thereof; R⁵ is C₂-C₄ linear alkylene, C₃-C₄ branched alkylene, and mixtures thereof; R⁶ is C₂-C₁₂ linear alkylene, and mixtures thereof; R⁷ is hydrogen, C₁-C₄ alkyl, and mixtures thereof; M is hydrogen or a water soluble cation; i is 0 or 1; i' is 0 or 1; j is 0 or 1; j' is 0 or 1; k is from 1 to 20; and m is from 0 to 10.

5. A composition according to Claim 4 wherein i and j are each equal to 0.

6. A composition according to Claim 1 wherein said zwitterionic polymeric suds stabilizer has the formula:




wherein R is C₁-C₁₂ linear alkylene, C₁-C₁₂ branched alkylene, and mixtures thereof; R¹ is a unit capable of having a negative charge at a pH of from 4 to 12; R² is a unit capable of having a positive charge at a pH of from 4 to 12; C₁-C₁₂ linear alkylene amino alkylene having the formula:




L¹, and mixtures thereof; wherein each R¹³ is independently L¹, ethylene, and mixtures thereof; each S is independently selected from C₁-C₁₂ linear alkylene, C₁-C₁₂ branched alkylene, C₃-C₁₂ linear alkenylene, C₃-C₁₂ branched alkenylene, C₃-C₁₂ hydroxyalkylene, C₄-C₁₂ dihydroxyalkylene, C₆-C₁₀ arylene, C₈-C₁₂ dialkylarylene, $-(R^5O)_kR^5-$, $-(R^5O)_kR^6(OR^5)_k-$, $-CH_2CH(OR^7)CH_2-$,

and mixtures thereof; L^1 is a linking unit independently selected from the following:




and mixtures thereof; $n^1 + n^2$ has a value such that said zwitterionic polymers suds stabilizer has an average molecular weight of from 1,000 to 2,000,000 daltons; n' is equal to n'' and further $n' + n''$ is less than or equal to 5% or the value $n^1 + n^2$; x is 0 to 6; y is 0 or 1; and z is 0 or 1.

7. A composition according to any of Claims 1-6 further comprising from 0.25% to 15% of a diamine wherein said diamine has the formula:




wherein each R^{20} is independently selected from the group consisting of hydrogen, C_1 - C_4 linear or branched alkyl, alkyleneoxy having the formula:



wherein R^{21} is C_2 - C_4 linear or branched alkylene, and mixtures thereof; R^{22} is hydrogen, C_1 - C_4 alkyl, and mixtures thereof; y is from 1 to 10; X is a unit selected from:

- i) C_3 - C_{10} linear alkylene, C_3 - C_{10} branched alkylene, C_3 - C_{10} cyclic alkylene, C_3 - C_{10} branched cyclic alkylene, an alkyleneoxyalkylene having the formula:



- wherein R^{21} and y are the same as defined herein above;
- ii) C_3 - C_{10} linear, C_3 - C_{10} branched linear, C_3 - C_{10} cyclic, C_3 - C_{10} branched cyclic alkylene, C_6 - C_{10} arylene, wherein said unit comprises one or more electron donating or electron withdrawing moieties which provide said diamine with a pK_a greater than 8; and
- iii) mixtures of (i) and (ii)
- provided said diamine has a pK_a of at least 8.
8. A composition according to Claim 7, wherein said diamine is 1,3-bis(methylamine)-cyclohexane.
9. A composition according to any of Claims 1-8 further comprising an enzyme selected from the group consisting of protease, amylase, and mixtures thereof.
10. A method for providing extended suds volume and suds duration when dishware in need of cleaning is washed, comprising the step of contacting said dishware with an aqueous solution of a liquid detergent comprising:
- a) an effective amount of a zwitterionic polymeric suds stabilizer;
 - b) an effective amount of a deterative surfactant; and
 - c) the balance carriers and other adjunct ingredients;
- provided the pH of a 10% aqueous solution of said composition is from 4 to 12.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/24699

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C11D3/37

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	GB 2 104 091 A (KAO CORP) 2 March 1983 see page 1, line 21 - page 3, line 2 see page 3, line 42 - line 45; examples 1,2; table 1 ---	1-5,10 6
X	EP 0 357 976 A (DOW CORNING) 14 March 1990 see abstract ---	1
A	WO 97 02337 A (PROCTER & GAMBLE) 23 January 1997 see page 16, paragraph 1 - page 17, paragraph 2 see page 20, paragraph 3 - paragraph 4; claim 1 ---	1,9,10
A	EP 0 232 092 A (GOLDMAN ROBERT ET AL) 12 August 1987 see claim 1; example 5 ---	1,7
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

16 March 1999

Date of mailing of the international search report

30/03/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentkan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Saunders, T

INTERNATIONAL SEARCH REPORT

Int .tional Application No

PCT/US 98/24699

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	<p>WO 98 28393 A (PROCTER & GAMBLE)</p> <p>2 July 1998</p> <p>see claims</p> <p>-----</p>	1,7-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/24699

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2104091 A	02-03-1983	JP 1055320 B	24-11-1989
		JP 1569529 C	10-07-1990
		JP 58013700 A	26-01-1983
		HK 46788 A	01-07-1988
EP 0357976 A	14-03-1990	US 4879051 A	07-11-1989
		AU 615097 B	19-09-1991
		AU 3937389 A	08-02-1990
		CA 1331121 A	02-08-1994
		DE 68923501 D	24-08-1995
		DE 68923501 T	01-02-1996
		JP 2068358 A	07-03-1990
		JP 2513851 B	03-07-1996
WO 9702337 A	23-01-1997	KR 9612273 B	18-09-1996
		NONE	
EP 0232092 A	12-08-1987	AU 590043 B	26-10-1989
		AU 6802687 A	30-07-1987
		JP 1062486 A	08-03-1989
WO 9828393 A	02-07-1998	AU 5379598 A	17-07-1998

Form PCT/ISA/210 (patent family annex) (July 1992)

